

PESTICIDE STANDARD WRITTEN NOTIFICATION

FOR SCHOOLS, DAY CARE PROGRAMS, AND SCHOOL-AGE CHILDCARE PROGRAMS

- The school, day care center, and/or school-age childcare program is responsible for sending this standard written notification form to employees, pupils, parents etc. to insure that they receive this information at least 2 working days prior to any pesticide use.
- It is recommended that the Pest Management Professional use this ready-to-copy standard written notification form for the purpose of providing pesticide use information to the school, day care center, and/or school-age childcare program. The Pest Management Professional should save this form for copying.

School: Franklin Schools
Name of School, Day care center, and/or School age childcare program

Pest Management Company: Natural Tree and Lawn Care P.O. Box 175 Avon MA 02322
(Please Print) Name Address

Pest Management Professional: Lawrence Taylor 0037746
(Please Print) License number
Mike Meglia Olink 0046566

A. List the Approximate Dates on which the pesticide use shall commence and conclude

Beginning Date April 14th 2019 **Ending Date** April 15th 2019

B. Record the specific location of the anticipated pesticide use

Athletic Fields, Keller Sullivan 500 Lincoln St., Remington Jefferson 628
Washington St., Davis Thayer 138 West Central St., Parmenter School 235
Wachusett St. Kennedy School 551 Pond St, High school 218 Oak St., Horace Mann
School 224 Oak St.

C. Pesticide Information (Pest Management Professional should be specific as is possible when listing product(s) to be used)

Pesticide Product Name	Pesticide Type	EPA Registration #	Description/Purpose of treatment and/or application
1. <u>Shaws 30-0-10</u>		<u>8378-73</u>	<u>Weed Control + Grub Control + Fertilizer</u>
2. <u>0.067% Acelepryn</u>	<u>Insecticide</u>		
3. <u>plus 0.222% Dimension</u>	<u>Herbicide</u>		
4. <u>Turf Care</u>		<u>100-1492-</u>	<u>Fertilizer with Grub Control</u>
5. <u>Grow Star 22-0-5</u>	<u>Insecticide</u>	<u>82757</u>	


This standard written notification must be accompanied by the following 2 documents. These materials are available from the DAR web page www.mass.gov/agr. Follow the links to the Children's Protection page.

- Chemical Specific Fact Sheet(s)
- Consumer Information Bulletin for school, day care center, and/or school-age childcare program.



Commonwealth of Massachusetts
Department of Agricultural Resources
Pesticide Bureau

251 Causeway St. Boston, MA 02114
Phone: 617-626-1781 Fax: 617-626-1850 Web:
<http://www.mass.gov/agr/>



**THE ACT PROTECTING CHILDREN AND FAMILIES
FROM HARMFUL PESTICIDES OF 2000**

Massachusetts Pesticide Bureau Consumer Information Bulletin FOR SCHOOLS, DAYCARE
CENTERS AND SCHOOL AGE CHILD CARE PROGRAMS

The Massachusetts Pesticide Control Act requires parents, staff, and children to receive this Massachusetts Pesticide Bureau Consumer Information bulletin whenever pesticide applications are being made on the property of your school, daycare center or school age child care program. This bulletin is being provided to you along with a Standard Written Notification form and a Pesticide Specific Factsheet.

Why am I receiving this information and what should I do when I receive it?

The purpose of the Standard Written Notification is to provide you with information about pesticide applications which are taking place on the property of your school, day care center or school age child care program. The bulletin provides information about precautions you can take to minimize exposure to any pesticides. The Pesticide Specific Factsheets provide information about the properties of the pesticides being used.

Who applies pesticides in my school, daycare center or school age child care program?

Commercial pest management professionals, facilities managers, grounds personnel or custodians. Regardless of the approach used, the person who applies the pesticides must have a current and valid Pesticide Bureau Applicator license. Check the standard written notification form for the applicator's license number.

How do I know when pesticides are being applied?

Employees, supervised children and their guardians must receive standard written notification at least two working days prior to the application of pesticides outdoors on the property. The standard written notification form, which accompanies this bulletin, includes:

- approximate dates when the application shall commence and conclude;
- specific location of the application;
- product name, type and EPA Registration number of the pesticide;
- a Pesticide Specific Fact Sheet;
- a description of the purpose of the application and
- this Consumer Information Bulletin

The notification must also be posted in a common area of the facility at least two working days before the outdoor application is to commence and at least 72 hours after the application. Treated areas will be posted with clear and conspicuous warning signs along the perimeter. This information will be supplied to the school by the licensed pesticide applicator.

Are applications of pesticides safe?

All pesticides must be treated with caution. They are intended to be specifically poisonous to target pest insects, weeds, mold, fungus etc. - and may also be harmful to other living things including humans.

Some degree of risk is always posed by their use. Because of this inherent risk, a number of regulatory and non-regulatory mechanisms have evolved to deal with those risks. Included among these mechanisms are pesticide regulations such as those enforced by the Massachusetts Pesticide Bureau; licensing and training of pesticide applicators; improved pesticide application methods; and the use of Integrated Pest Management (IPM).

What precautions can I take to minimize my exposure to pesticide applications?

There are several precautions that can be taken to reduce potential exposure to pesticides. These precautions will vary depending on where and how the pesticides are applied. Chemicals may be ingested, inhaled and absorbed through the skin. Know where the pesticide will be applied and how you might come into contact with it. Use common sense. The licensed pesticide applicator is required to post yellow signs to indicate a pesticide application on school grounds. These are some suggested general precautions. Ask the licensed pesticide applicator for other suggestions or directions specific to the work being done.

For outdoor applications:

- be familiar with the small yellow signs which applicators are required to post when a pesticide is applied outdoors to turf. Stay off the field until the flags are removed.
- if you are sensitive to chemicals, avoid the area of pesticide application for 72 hours.
- ensure that pets are kept away from the area of pesticide application

For indoor applications:

- cover or refrigerate edible products.
- remove or cover toys, clothes, and bedding from areas to be treated.
- remove pets including their food and water bowls and toys from the area to be treated.
- ventilate as much as possible during and, following an indoor pesticide application, open the windows.
- do not walk on treated areas and carpets until completely dry. Ask about drying times.

What types of pesticides will be applied?

Pesticide applicators may apply pesticides in several forms for control of insects and weeds. Dusts, aerosol sprays, sprays, baits, and fogs are all common forms in which pesticides exist and are used. For control of termites, the soil around the building may be impregnated with a pesticide. To control weeds, pesticides may be used as granules or sprays. Mechanical traps may also be used to control rodents.

In Massachusetts schools daycare centers and school age child care programs have to develop special pest management plans called Integrated Pest Management (IPM) plans. IPM is an approach to pest management which relies on a combination of common sense practices, including pesticides, for preventing and controlling pests. All plans are required to be submitted to the Department of Agricultural Resources. Check the DAR website to see if your school has submitted its plan.

<http://massnrc.org/ipm/index.html>

What if I have a question or problem?

Questions about what pesticides will be applied and why, and specific information about the application should be referred to the licensed pesticide applicator doing the work.

The Massachusetts Department of Agricultural Resources, Pesticide Bureau is responsible for enforcing the pesticide regulations and laws. Contact the Pesticide Bureau at 617-626-1781. Additional information can be found at the Pesticide Bureau website: <http://www.mass.gov/agr/pesticides/>

Updated April 2008.

**Shaw's® Turf Food with 0.067%
Acelepryn™ Plus 0.222% Dimension®**
INTENDED FOR USE BY COMMERCIAL APPLICATORS
ONLY.

For control of white grubs, other listed insect pests, and listed weeds in established lawns and ornamental turfgrass (including golf courses).

Not for Sale, Sale into, Distribution and/or Use in Nassau, Suffolk, Kings, Queens Counties of New York State.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Active Ingredients	By Weight
Chlorantranilprole*	0.067%
3-Bromo-N-[4-chloro-2-methyl-6- [(methylamino)carbonyl] phenyl]-1-(3-chloro- 2-pyridinyl)-1H-pyrazole-5-carboxamide	
Dithiopyr**	0.222%
3,5- pyridinedicarbothioic acid, 2- (difluoromethyl)- 4-(2-methylpropyl)-6- (trifluoromethyl)-S,S-dimethyl ester	
Other Ingredients	99.711%
TOTAL	100.000%
EPA Reg. No. 8378-73	EPA Est. No. 8378-IN-1

Shaw's® is a registered trademark of Knox Fertilizer Company, Inc. Acelepryn™ is a trademark of E.I. DuPont de Nemours and Company or its affiliates. Acelepryn™ is a trademark or registered trademark of DuPont used under license by Knox Fertilizer Company, Inc. Dimension® is a registered trademark of Dow AgroSciences LLC.

30-0-10

**FAIRWAY
GUARANTEED ANALYSIS :**

TOTAL NITROGEN (N) 30%

29.98% UREA NITROGEN*

0.02% WATER SOLUBLE NITROGEN

SOLUBLE POTASH (K₂O) 10%

SULFUR (S) 3.00%

3.00% COMBINED SULFUR (S)

DERIVED FROM: POLYMER-ENCAPSULATED UREA,
AMINO ACIDS AND SULFATE OF POTASH.

*39.98% SLOWLY AVAILABLE NITROGEN FROM:
POLYMER-ENCAPSULATED UREA.



GRO-GARD® (A proprietary blend of amino acids)

KNOX FERTILIZER COMPANY, INC.

P.O. BOX 248

KNOX, IN 46534

NET WEIGHT 50 LBS. (22.7 KG.)

5329289

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in Eyes: hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the container or label with you when calling a poison control center or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC
ANIMALS**

CAUTION! Causes moderate eye irritation. Avoid contact with eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Shoes plus socks.

After the product has been loaded into properly calibrated application equipment, shirt, pants, socks, and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Do not apply directly to water. Do not apply where/when conditions favor runoff. Runoff from treated turf may adversely affect aquatic organisms in water adjacent to use sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory:

This product may contaminate water through runoff. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water table are more prone to produce runoff that contains this product. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Fall Application: When this product is applied in the fall, a sequential or second application may be made the following spring to provide up to 4 months control of weeds. If the product is applied in the fall, do not reseed, overseed or sprig turfgrasses before 12 weeks after application due to potential injury to newly reseeded, overseeded or sprigged turfgrasses.

Weeds Controlled: This product will provide control of crabgrass and other grass and broadleaf weeds listed below when applied prior to germination in established lawns and ornamental grasses.

Grass Weeds	Broadleaf Weeds
Annual Bluegrass	Bittercress
Annual Ryegrass	Buttercup Oxalis
Barnyardgrass	Carolina Geranium
Crabgrass	Chickweed
Crowfootgrass	Common Lespedeza
Goosegrass	Common Purslane
Green Foxtail	Corn Speedwell
Kikuyugrass (seedling)	Creeping Oxalis
Perennial Ryegrass	Henbit
Sandbur	Mustard
Smutgrass (seedling)	Parsley-piert
Yellow Foxtail	Pineappleweed
	Prostrate Spurge
	Shepardspurse
	Spotted Spurge
	Yellow Woodsorrel

APPLICATION RATES FOR LAWNS, GOLF COURSES AND OTHER RECREATIONAL TURFGRASS AREAS

Table 1 provides the application rates that will control the listed insect pests and will also control the listed grass and broadleaf weeds. **Restrictions:** Do not apply more than 225 pounds of this product/acre per application and no more than 338 pounds (equivalent to 0.23 lb of chlorantraniliprole and 0.75 lb of dithiopyr) of product/acre per year in broadcast applications to turfgrass.

TABLE 1: TURFGRASS APPLICATION RATES

Target Pest (Including Listed Weeds)	Product per Acre	Product per 1,000 Sq. Ft.	Lb A/A Acelepryn	Lb A/A Dimension
White Grubs (including <i>Aphodius</i> spp., Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, May/June beetles (<i>Phyllophaga</i> spp.), northern masked chafer, oriental beetle and southern masked chafer)	150 to 225 pounds	3.44 to 5.16 pounds	0.1 to 0.15	0.33 to 0.50
Billbugs				
European Crane Fly				
Turf caterpillars (including armyworms, cutworms and sod webworms)				

NOTE: For 3-4 months weed control use 150 pounds/acre of this product.

For early postemergence control of crabgrass, use 150 to 225 pounds of this product/acre. Use the higher rate in areas of high weed pressure.

In the State of New York: Do not exceed 0.5 pounds of dithiopyr/A (225 pounds product/A) per year.

SPECIFIED SPREADER SETTINGS

Spreader	3.44	5.16
Scott's R8A	L - M	N - O
Lesco High Wheel	G - H	I - J
Spyker 24-22	5 ¼ - 5 ½	5 ½ - 6
Prizelawn	J - K	L - M
Earthway	14 - 15	16 - 17
Vicon	20	24
Lely	4 ½	5 1/2

CLEANING APPLICATION EQUIPMENT

Thoroughly clean all application equipment after use to reduce the risk of forming hardened deposits that might become difficult to remove. Thoroughly rinse application equipment with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean equipment near wells, or other water sources. Dispose of waste rinse water in accordance with local regulations.

INTEGRATED PEST MANAGEMENT (IPM) PROGRAMS

Shaw's® Turf Food with 0.067% Acelepryn™ Plus 0.222% Dimension is recommended for IPM programs on turf because it does not directly impact natural arthropod predator and parasitoid populations including ladybird beetles, lacewings, minute pirate bugs and predatory mites. The feeding behavior of beneficial arthropods will aid in extending natural control of other insect and mite pests and will reduce the possibility of secondary pest outbreaks. This product will reduce the target pest species that may serve as a food source for beneficial arthropods.

RESISTANCE MANAGEMENT

Some insects are known to develop resistance to products used repeatedly for control. When this occurs, the labeled application rates fail to suppress the pest population below threshold levels. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of active classes of insecticides on succeeding generations and targeting the most susceptible life stage. Consult your local Knox Fertilizer Company representative, Cooperative Extension Service specialist or pest control advisor for the latest information on resistance management in your area.

USE PRECAUTIONS & RESTRICTIONS

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production.
- Do not formulate this product into other end-use products.
- Do not apply this product in commercial nurseries or greenhouses.
- Do not apply this product on golf course putting greens.
- Apply only to lawns and turfgrasses that are well established. Do not apply to lawns or turfgrasses that are under stress.
- Do not apply more than 225 pounds of product/A (equivalent to 0.15 lb of chlorantraniliprole/A and 0.5 lb of dithiopyr/A) per application and no more than 338 pounds of product/A (equivalent to 0.23 lb of

Ground Water Advisory:

This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label before using this product.

Shaw's® Turf Food with 0.067% Acelepryn™ Plus 0.222% Dimension® must be used only in accordance with specifications on this label or in separate Knox Fertilizer Company supplemental labeling that may be made temporarily available through local distributors as a result of new EPA approvals. Knox Fertilizer Company is not responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by Knox Fertilizer Company. User assumes all risks associated with such non-specified use.

PRODUCT INFORMATION

Shaw's® Turf Food with 0.067% Acelepryn™ Plus 0.222% Dimension® is a combination product containing an Insecticide (Acelepryn™) and a herbicide (Dimension®) and may be applied for the control of white grubs, other listed insect pests, and listed weeds infesting recreational turfgrass (including golf courses).

This product may be used on established turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, commercial lawns, industrial facilities, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, parks, playgrounds, schools, day-care facilities, golf courses (tee box areas, roughs, fairways, collars etc.), athletic fields, and other landscaped areas.

This product must not be applied when the soil is saturated with water because adequate distribution of the active ingredients vertically in the soil profile cannot be achieved under this condition.

The insecticide component of this product (Acelepryn™) is in GROUP 28 of the EPA's Insecticide and acaricide Groups Based on Target Site of Action (EPA PR Notice 2001-5) and this product may be used in rotational resistance management programs. Consult your local Knox Fertilizer Company representative, Cooperative Extension Service specialist or pest control advisor for regionally specific information regarding application timing.

TOLERANT TURFGRASS SPECIES

This product may be applied to established lawns and ornamental turfgrasses composed of the following turfgrass species:

Cool-Season Turfgrass	Warm-Season Turfgrass
Creeping Bentgrass	Bahiagrass
Fine Fescues	Bermudagrass
Kentucky Bluegrass	Buffalograss
Perennial Ryegrass	Carpetgrass
Tall Fescues	Centipedegrass
	St. Augustinegrass
	Zoysiagrass

APPLICATION EQUIPMENT

Apply Shaw's® Turf Food with 0.067% Acelepryn™ Plus 0.222% Dimension® with rotary-type equipment. The application equipment must be properly calibrated according to the manufacturer's directions to ensure uniform coverage over the treated area. During application, check the equipment periodically to confirm that it is working properly and that the granules are being uniformly distributed. A more uniform application can be made by spreading one-half of the recommended amount of this product over the turfgrass area and then spreading the remaining one-half of this product at a right angle to the previous application direction.

APPLICATION TIMING – INSECT CONTROL

White Grubs: Apply this product from early April to early September for preventative and early curative control of all of the major white grub species infesting turfgrass. Irrigate turf immediately after application.

Billbugs: Apply this product when overwintered adult billbugs are first observed. This will usually occur in late April or early May in regions with cool-season turfgrasses. An application of this product at this time will also provide white grub control. The higher rates listed in Table 1 may be required for applications made after early May or when billbug species other than bluegrass billbug are present.

European Crane Fly: Apply this product between September and November to control European crane fly larvae in turfgrass. The higher rate listed in Table 1 may be required to achieve control when applications are made in November.

Turf Caterpillars: This product will provide curative caterpillar control in turfgrass. If the area being treated is maintained at a mowing height of greater than one inch, the higher rate listed in Table 1 may be required during periods of high pest pressure.

APPLICATION TIMING – WEED CONTROL

This product may be applied in the spring or fall prior to weed seed germination in established lawns and ornamental turfgrasses. This product provides preemergence control of crabgrass and other weeds when applied prior to their germination, and it may also provide postemergence control of seedling crabgrass.

Postemergence Crabgrass Control: The postemergence control provided by this product is limited to crabgrass, and it will only provide control up to 4 weeks after crabgrass germination, with best control expected when crabgrass is small (3-4 leaf stage). This product will not control large, tillered crabgrass. For control of other grasses and broadleaf weeds, this product must be applied as a preventative treatment before their emergence.

Reseeding, Overseeding or Sprigging: Reseeding, overseeding, or sprigging of turfgrasses listed on this label into areas that have been previously treated with this product must be delayed until 8 weeks after the previous application. Reseeding, overseeding, or sprigging before 8 weeks after application may injure or prevent the establishment of the newly reseeded, overseeded, or sprigged turfgrass.

Use on New Lawns of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue: To establish a new lawn of Kentucky Bluegrass, Perennial Ryegrass and/or Tall Fescue, this product may be applied 2 weeks after the first sign of germination. For other tolerant turfgrasses listed on this label, wait until the grass is well established (2 mowings) before applying this product.

(OVER)



Acelepryn® 0.067%

On TCS GrowStar® Fertilizer

GROUP 28 INSECTICIDE

INTENDED FOR USE BY COMMERCIAL APPLICATORS ONLY.

For systemic control of white grubs and other listed pests infesting landscape and recreational turfgrass (including golf courses) as well as landscape ornamentals, interior plantscapes and sod farms.

Not for Sale, Sale Into, Distribution and/or Use In Nassau, Suffolk, Kings, Queens Counties of New York State.

ACTIVE INGREDIENTS:	By Weight
Chlorantraniliprole* 3-Bromo-N-[4-chloro-2-methyl-6-[(methylamino)carbonyl]phenyl]-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide.....	0.067%
Other Ingredients.....	99.933%
Total.....	100.000%

* Chlorantraniliprole belongs to the anthranilic diamide chemical class.

22-0-5

GUARANTEED ANALYSIS

AVAILABLE Nitrogen (N).....22.0%
22.0% Urea Nitrogen*

SOLUBLE POTASH (K₂O).....5.0%

DERIVED FROM: Polymer Sulfur-Coated Urea, Urea, Muriate of Potash.

CHLORINE (Cl) Max.....3.8%

*8.1% Slowly Available Nitrogen from Polymer Sulfur-Coated Urea.

In the state of FL it is recommend that the user of this product follow the Florida Green Industries BMPs at: http://www.flaes.org/pdf/BMP_Book_final.pdf
In the state of FL it is recommend that the user of this product follow the Florida Golf Course BMPs at: <http://www.flaes.org/pdf/gltbmp07.pdf>

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
------------	---

Have the container or label with you when calling a poison center or doctor, or going for treatment. You may also contact 1-800-441-3637 emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks.

After the product has been loaded into properly calibrated application equipment, shirt, pants, socks and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instruction for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites.

Surface Water Advisory: This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory: This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label before using this product.

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer must be used only in accordance with specifications on this label or in separate Turf Care Supply Corp. supplemental labeling that may be made temporarily available through local distributors, as a result of new EPA approvals. Turf Care Supply Corp. will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by Turf Care Supply Corp. User assumes all risks associated with such non-specified use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with it labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, residential, industrial and commercial lawns and sports fields are not within the scope of the Worker Protection Standard.

Do not allow other workers to enter the treated area while application is in progress.

PRODUCT INFORMATION

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer may be applied for the control of white grubs and other listed pests infesting landscape and recreational turfgrass (including golf courses) as well as landscape ornamentals and interior plantscapes.

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer may be used on plants or turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, commercial lawns, industrial facilities, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes.

ACTIVE INGREDIENTS:

Chlorantraniliprole* 3-Bromo-N-[4-chloro-2-methyl-6-[(methylamino)carbonyl]phenyl]-1-(3-chloro-2-pyridinyl)-1H-pyrazole-5-carboxamide.....	0.067%
Other Ingredients.....	99.933%
Total.....	100.000%

By Weight

* Chlorantraniliprole belongs to the anthranilic diamide chemical class.

22-0-5

GUARANTEED ANALYSIS

AVAILABLE Nitrogen (N).....22.0%
22.0% Urea Nitrogen*

SOLUBLE POTASH (K₂O).....5.0%

DERIVED FROM: Polymer Sulfur-Coated Urea, Urea, Muriate of Potash.

CHLORINE (Cl) Max.....3.8%

*8.1% Slowly Available Nitrogen from Polymer Sulfur-Coated Urea.

In the state of FL it is recommend that the user of this product follow the Florida Green Industries BMPs at: http://www.flaes.org/pdf/BMP_Book_final.pdf

In the state of FL it is recommend that the user of this product follow the Florida golf Course BMPs at: <http://www.flaes.org/pdf/glfbmp07.pdf>

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
------------	---

Have the container or label with you when calling a poison center or doctor, or going for treatment. You may also contact 1-800-441-3637 emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution! Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks.

After the product has been loaded into properly calibrated application equipment, shirt, pants, socks and shoes are sufficient Personal Protective Equipment. Follow manufacturer's instruction for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

EPA REG. NO. 100-1492-82757

EPA EST. NO. 82757-OH-001 (MF) First letter of lot code indicates manufacturing site.

EPA EST. NO. 82757-MA-002 (H) First letter of lot code indicates manufacturing site.

EPA EST. NO. 82757-FL-001 (S) First letter of lot code indicates manufacturing site.

Net Contents 50 lbs. (22.67 kg)

reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory: This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label before using this product.

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer must be used only in accordance with specifications on this label or in separate Turf Care Supply Corp. supplemental labeling that may be made temporarily available through local distributors, as a result of new EPA approvals. Turf Care Supply Corp. will not be responsible for losses or damages resulting from the use of this product in any manner not specifically stated on this label or other labels or bulletins published by Turf Care Supply Corp. User assumes all risks associated with such non-specified use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with it labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about personal protective equipment, restricted-entry interval, and notification to workers (as applicable).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Professional applications to golf courses, residential, industrial and commercial lawns and sports fields are not within the scope of the Worker Protection Standard.

Do not allow other workers to enter the treated area while application is in progress.

PRODUCT INFORMATION

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer may be applied for the control of white grubs and other listed pests infesting landscape and recreational turfgrass (including golf courses) as well as landscape ornamentals and interior plantscapes.

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer may be used on plants or turfgrasses that are being grown for aesthetic or recreational purposes or climatic modification in or around home lawns, commercial lawns, industrial facilities, residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, interior plantscapes, ornamental gardens, parks, wildlife plantings, playgrounds, schools, day-care facilities, golf courses (tee box areas, roughs, fairways, greens, collars etc.), athletic fields, sod farms and other landscaped areas.

Do not apply ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer when the soil is saturated with water because adequate distribution of the active ingredient vertically in the soil profile cannot be achieved under this condition.

ACELEPRYN® 0.067% on TCS GrowStar® Fertilizer is in GROUP 28 of the EPA's Insecticide and Acaricide Groups Based on Target Site of Action (EPA PR Notice 2001-5) and may be used in rotational resistance management programs.

Consult your local Turf Care Supply Corp. representative, Cooperative Extension Service specialist or pest control advisor for regionally specific information regarding application timing.

#901844

